

---

---

# Generalized matrix $t$ distribution based on new matrix gamma distribution

---

---

Authors: ANIS IRANMANESH\*

- Department of Mathematics and Statistics, Mashhad Branch, Islamic Azad University, Mashhad, Iran.  
(iranmanesh0030@mshdiau.ac.ir)

DAYA K. NAGAR

- Instituto de Matemáticas, Universidad de Antioquia, Medellín Colombia.  
(dayaknagar@yahoo.com)

SARA SHOKRI

- Department of Mathematics and Statistics, Mashhad Branch, Islamic Azad University, Mashhad, Iran.  
(sara.shokri30@gmail.com)

SARALEES NADARAJAH

- School of Mathematics, University of Manchester, United Kingdom.  
(Saralees.Nadarajah@manchester.ac.uk)

Abstract:

- In this paper a generalized matrix variate gamma distribution, which includes a trace function in the kernel of the density, is introduced. Some important statistical properties including Laplace transform, distributions of functions, expected value of the determinant and expected value of zonal polynomial of the generalized gamma matrix are derived. Further, by using the distribution of the inverse of this newly defined generalized gamma matrix as the prior for the characteristic matrix of a matrix variate normal distribution, a new generalized matrix  $t$  type family of distributions is generated. Some important statistical characteristics of this family are also exhibited.

Key-Words:

- *Bayes estimator, Generalized matrix gamma distribution, Generalized matrix  $t$  distribution, Zonal polynomial.*

AMS Subject Classification:

- 62E05; 62H99.



---

## 1. INTRODUCTION

---

In 1964, Lukacs and Laha defined the matrix variate gamma (MG) distribution. In multivariate statistical analysis, the MG distribution has been the subject of considerable interest, study, and applications for many years. For example, the Wishart distribution, which is the distribution of the sample variance covariance matrix when sampling from a multivariate normal distribution, is a special case of the MG distribution. Applications of the MG distribution have included: damping modeling (Adhikari [1]); models for stochastic upscaling for inelastic material behavior from limited experimental data (Das and Ghanem [7], [8]); models for fusion yield [15]; models for uncertainty quantification (Pascual and Adhikari [31]); characterizing the distribution of anisotropic micro-structural environments with diffusion-weighted imaging (Scherrer et al. [32]); models for magnetic tractography (Chamberland et al. [4]); models for diffusion compartment imaging (Scherrer et al. [33]); models for image classification (Luo et al. [24]); models for accurate signal reconstruction (Jian et al. [22], Bates et al. [3]). Two recent applications of the Wishart distribution can be found in Arashi et al. [2] and Ferreira et al. [13].

However, generalizations of the MG distribution have been neglected and there is no account on this matter in the literature. The only extension that we are aware of is the inverted matrix variate gamma distribution due to Iranmanesh et al. [17]: if  $\mathbf{X}$  has the MG distribution then  $\mathbf{X}^{-1}$  has the inverted matrix variate gamma distribution. A generalization of the MG distribution must contain the MG distribution as a particular case. See also Iranmanesh et al. [18] and references there in for more details.

The goal of this paper is to give the first generalization to the MG distribution, where its kernel includes zonal polynomials (Takemura [34]). The generalization proposed has two shape parameters. One of the shape parameters acts on the determinant of the data while the other acts on the trace of the data. The MG distribution has only one shape parameter acting on the determinant of the data. The proposed generalization can be more flexible for data modeling if: i) both trace and determinant are significant (that is, the empirical distribution of the data has significant patterns involving both the trace and determinant); ii) if trace is significant but determinant is not (that is, the empirical distribution of the data has significant patterns involving only the trace); iii) if trace is more significant than determinant is (that is, the empirical distribution of the data has more significant patterns involving the trace).

For our purpose, we first provide the reader with some preliminary definitions and lemmas. Most of the following definitions and results can be found in Gupta and Nagar [14], Muirhead [27], and Mathai [26].

---

## 2. Preliminaries

---

In this section we state certain well known definitions and results. These results will be used in subsequent sections.

Let  $\mathbf{A} = (a_{ij})$  be a  $p \times p$  matrix. Then,  $\mathbf{A}'$  denotes the transpose of  $\mathbf{A}$ ;  $\text{tr}(\mathbf{A}) = a_{11} + \cdots + a_{pp}$ ;  $(\mathbf{A}) = \exp(\text{tr}(\mathbf{A}))$ ;  $\det(\mathbf{A}) =$  determinant of  $\mathbf{A}$ ; norm of  $\mathbf{A} = \|\mathbf{A}\| =$  maximum of absolute values of eigenvalues of the matrix  $\mathbf{A}$ ;  $\mathbf{A}^{1/2}$  denotes a symmetric positive definite square root of  $\mathbf{A}$ ;  $\mathbf{A} > 0$  means that  $\mathbf{A}$  is symmetric positive definite and  $0 < \mathbf{A} < \mathbf{I}_p$  means that the matrices  $\mathbf{A}$  and  $\mathbf{I}_p - \mathbf{A}$  are symmetric positive definite. The multivariate gamma function which is frequently used in multivariate statistical analysis is defined by

$$\begin{aligned} \Gamma_p(a) &= \int_{\mathbf{X} > 0} (-\mathbf{X}) \det(\mathbf{X})^{a-(p+1)/2} d\mathbf{X} \\ &= \pi^{p(p-1)/4} \prod_{i=1}^p \Gamma(a - \frac{i-1}{2}), \quad \text{Re}(a) > \frac{p-1}{2}. \end{aligned} \quad (2.1)$$

Let  $C_\kappa(\mathbf{X})$  be the zonal polynomial of  $p \times p$  complex symmetric matrix  $\mathbf{X}$  corresponding to the ordered partition  $\kappa = (k_1, \dots, k_p)$ ,  $k_1 \geq \cdots \geq k_p \geq 0$ ,  $k_1 + \cdots + k_p = k$  and  $\sum_\kappa$  denotes summation over all partitions  $\kappa$  of  $k$ . The generalized hypergeometric coefficient  $(a)_\kappa$  used above is defined by

$$(a)_\kappa = \prod_{i=1}^p (a - \frac{i-1}{2})_{k_i},$$

where  $(a)_r = a(a+1) \cdots (a+r-1)$ ,  $r = 1, 2, \dots$  with  $(a)_0 = 1$ .

**Lemma 2.1.** *Let  $\mathbf{Z}$  be a complex symmetric  $p \times p$  matrix with  $\text{Re}(\mathbf{Z}) > 0$ , and let  $\mathbf{Y}$  be a symmetric  $p \times p$  matrix. Then, for  $\text{Re}(a) > (p-1)/2$ , we have*

$$\int_{\mathbf{X} > 0} (-\mathbf{X}\mathbf{Z}) (\det \mathbf{X})^{a-(p+1)/2} C_\kappa(\mathbf{X}\mathbf{Y}) d\mathbf{X} = (a)_\kappa \Gamma_p(a) (\det \mathbf{Z})^{-a} C_\kappa(\mathbf{Y}\mathbf{Z}^{-1}). \quad (2.2)$$

**Lemma 2.2.** *Let  $\mathbf{Z}$  be a complex symmetric  $p \times p$  matrix with  $\text{Re}(\mathbf{Z}) > 0$ , and let  $\mathbf{Y}$  be a symmetric  $p \times p$  matrix. Then, for  $\text{Re} > (p-1)/2$ , we have*

$$\int_{\mathbf{X} > 0} (-\mathbf{X}\mathbf{Z}) (\det \mathbf{X})^{a-(p+1)/2} [\text{tr}(\mathbf{X}\mathbf{Y})]^k d\mathbf{X} = \Gamma_p(a) (\det \mathbf{Z})^{-a} \sum_{\kappa} (a)_\kappa C_\kappa(\mathbf{Y}\mathbf{Z}^{-1}). \quad (2.3)$$

For  $\mathbf{Z} = \mathbf{Y}$  in (2.3), we get

$$\begin{aligned} \int_{\mathbf{X} > 0} (-\mathbf{X}\mathbf{Y}) (\det \mathbf{X})^{a-(p+1)/2} [\text{tr}(\mathbf{X}\mathbf{Y})]^k d\mathbf{X} &= \Gamma_p(a) (\det \mathbf{Y})^{-a} \sum_{\kappa} (a)_\kappa C_\kappa(\mathbf{I}_p) \\ &= \Gamma_p(a) (ap)_k (\det \mathbf{Y})^{-a}. \end{aligned} \quad (2.4)$$

The above result was derived by Khatri [23].

Davis [9, 10] introduced a class of polynomials  $C_\phi^{\kappa,\lambda}(\mathbf{X}, \mathbf{Y})$  of  $p \times p$  symmetric matrix arguments  $\mathbf{X}$  and  $\mathbf{Y}$ , which are invariant under the transformation  $(\mathbf{X}, \mathbf{Y}) \rightarrow (\mathbf{H}\mathbf{X}\mathbf{H}', \mathbf{H}\mathbf{Y}\mathbf{H}')$ ,  $\mathbf{H} \in O(p)$ . For properties and applications of invariant polynomials we refer to Davis [9, 10], Chikuse [5] and Nagar and Gupta [28]. Let  $\kappa, \lambda, \phi$  and  $\rho$  be ordered partitions of non-negative integers  $k, \ell, f = k + \ell$  and  $r$ , respectively, into not more than  $p$  parts. Then

$$(2.5) \quad C_\phi^{\kappa,\lambda}(\mathbf{X}, \mathbf{X}) = \theta_\phi^{\kappa,\lambda} C_\phi(\mathbf{X}), \quad \theta_\phi^{\kappa,\lambda} = \frac{C_\phi^{\kappa,\lambda}(\mathbf{I}_p, \mathbf{I}_p)}{C_\phi(\mathbf{I}_p)},$$

$$(2.6) \quad C_\phi^{\kappa,\lambda}(\mathbf{X}, \mathbf{I}_p) = \theta_\phi^{\kappa,\lambda} \frac{C_\phi(\mathbf{I}_p) C_\kappa(\mathbf{X})}{C_\kappa(\mathbf{I}_p)}, \quad C_\phi^{\kappa,\lambda}(\mathbf{I}_p, \mathbf{Y}) = \theta_\phi^{\kappa,\lambda} \frac{C_\phi(\mathbf{I}_p) C_\lambda(\mathbf{Y})}{C_\lambda(\mathbf{I}_p)},$$

$$C_\kappa^{\kappa,0}(\mathbf{X}, \mathbf{Y}) \equiv C_\kappa(\mathbf{X}), \quad C_\lambda^{0,\lambda}(\mathbf{X}, \mathbf{Y}) \equiv C_\lambda(\mathbf{Y})$$

and

$$(2.7) \quad C_\kappa(\mathbf{X}) C_\lambda(\mathbf{Y}) = \sum_{\phi \in \kappa \cdot \lambda} \theta_\phi^{\kappa,\lambda} C_\phi^{\kappa,\lambda}(\mathbf{X}, \mathbf{Y}),$$

where  $\phi \in \kappa \cdot \lambda$  signifies that irreducible representation of  $Gl(p, R)$  indexed by  $2\phi$ , occurs in the decomposition of the Kronecker product  $2\kappa \otimes 2\lambda$  of the irreducible representations indexed by  $2\kappa$  and  $2\lambda$ . Further,

$$\begin{aligned} & \int_{\mathbf{R} > 0} (-\mathbf{C}\mathbf{R}) \det(\mathbf{R})^{t-(p+1)/2} C_\phi^{\kappa,\lambda}(\mathbf{A}\mathbf{R}\mathbf{A}', \mathbf{B}\mathbf{R}\mathbf{B}') d\mathbf{R} \\ &= \Gamma_p(t, \phi) \det(\mathbf{C})^{-t} C_\phi^{\kappa,\lambda}(\mathbf{A}\mathbf{C}^{-1}\mathbf{A}', \mathbf{B}\mathbf{C}^{-1}\mathbf{B}'), \end{aligned} \quad (2.8)$$

$$\begin{aligned} & \int_0^{\mathbf{I}_p} \det(\mathbf{R})^{t-(p+1)/2} \det(\mathbf{I}_p - \mathbf{R})^{u-(p+1)/2} C_\phi^{\kappa,\lambda}(\mathbf{R}, \mathbf{I}_p - \mathbf{R}) d\mathbf{R} \\ &= \frac{\Gamma_p(t, \kappa) \Gamma_p(u, \lambda)}{\Gamma_p(t+u, \phi)} \theta_\phi^{\kappa,\lambda} C_\phi(\mathbf{I}_p) \end{aligned} \quad (2.9)$$

and

$$\begin{aligned} & \int_0^{\mathbf{I}_p} \det(\mathbf{R})^{t-(p+1)/2} \det(\mathbf{I}_p - \mathbf{R})^{u-(p+1)/2} C_\phi^{\kappa,\lambda}(\mathbf{A}\mathbf{R}, \mathbf{B}\mathbf{R}) d\mathbf{R} \\ &= \frac{\Gamma_p(t, \phi) \Gamma_p(u)}{\Gamma_p(t+u, \phi)} C_\phi^{\kappa,\lambda}(\mathbf{A}, \mathbf{B}). \end{aligned} \quad (2.10)$$

In expressions (2.8), (2.9) and (2.10),  $\Gamma_p(a, \rho)$  is defined by

$$(2.11) \quad \Gamma_p(a, \rho) = (a)_\rho \Gamma_p(a).$$

Note that  $\Gamma_p(a, 0) = \Gamma_p(a)$ , which is the multivariate gamma function.

Let  $\mathbf{A}$ ,  $\mathbf{B}$ ,  $\mathbf{X}$  and  $\mathbf{Y}$  be  $p \times p$  symmetric matrices. Then

$$\int_{\mathbf{H} \in O(p)} C_{\phi}^{\kappa, \lambda}(\mathbf{A}\mathbf{H}'\mathbf{X}\mathbf{H}, \mathbf{B}\mathbf{H}'\mathbf{Y}\mathbf{H}) [d\mathbf{H}] = \frac{C_{\phi}^{\kappa, \lambda}(\mathbf{A}, \mathbf{B}) C_{\phi}^{\kappa, \lambda}(\mathbf{X}, \mathbf{Y})}{\theta_{\phi}^{\kappa, \lambda} C_{\phi}(\mathbf{I}_p)}, \quad (2.12)$$

where  $[d\mathbf{H}]$  is the unit invariant Haar measure. The above result is a generalization of Davis [10, Eq. 5.4] and is due to Díaz-García [11]. Finally, using (2.9) and (2.12), it is straightforward to see that

$$\begin{aligned} & \int_0^{\mathbf{I}_p} \det(\mathbf{R})^{t-(p+1)/2} \det(\mathbf{I}_p - \mathbf{R})^{u-(p+1)/2} C_{\phi}^{\kappa, \lambda}(\mathbf{A}\mathbf{R}, \mathbf{B}(\mathbf{I}_p - \mathbf{R})) d\mathbf{R} \\ &= \frac{\Gamma_p(t, \kappa) \Gamma_p(u, \lambda)}{\Gamma_p(t+u, \phi)} C_{\phi}^{\kappa, \lambda}(\mathbf{A}, \mathbf{B}). \end{aligned} \quad (2.13)$$

**Definition 2.1.** The  $n \times p$  random matrix  $\mathbf{X}$  is said to have a matrix variate normal distribution with  $n \times p$  mean matrix  $\mathbf{M}$  and  $np \times np$  covariance matrix  $\mathbf{\Omega} \otimes \mathbf{\Sigma}$ , denoted by  $\mathbf{X} \sim N_{n,p}(\mathbf{M}, \mathbf{\Omega} \otimes \mathbf{\Sigma})$ , if its probability density function (p.d.f) is given by (Gupta and Nagar [14])

$$\begin{aligned} & (2\pi)^{-np/2} \det(\mathbf{\Omega})^{-p/2} \det(\mathbf{\Sigma})^{-n/2} \\ & \times \exp \left\{ -\frac{1}{2} \text{tr} [\mathbf{\Omega}^{-1}(\mathbf{X} - \mathbf{M})\mathbf{\Sigma}^{-1}(\mathbf{X} - \mathbf{M})'] \right\}, \quad \mathbf{X} \in \mathbb{R}^{n \times p}, \quad \mathbf{M} \in \mathbb{R}^{n \times p}, \end{aligned}$$

where  $\mathbf{\Sigma}(p \times p) > 0$  and  $\mathbf{\Omega}(n \times n) > 0$ .

If  $\mathbf{X} \sim N_{n,p}(\mathbf{M}, \mathbf{\Omega} \otimes \mathbf{\Sigma})$ , then the characteristic function of  $\mathbf{X}$  is

$$\begin{aligned} \phi_{\mathbf{X}}(\mathbf{Z}) &= E[\exp(\text{tr}(\iota \mathbf{Z}' \mathbf{X}))] \\ &= \exp \left[ \text{tr} \left( \iota \mathbf{Z}' \mathbf{M} - \frac{1}{2} \mathbf{Z}' \mathbf{\Omega} \mathbf{Z} \mathbf{\Sigma} \right) \right], \quad \mathbf{Z} \in \mathbb{R}^{n \times p}, \quad \iota = \sqrt{-1}. \end{aligned}$$

The present paper has been organized in the following sections. In Section 3, a new generalized matrix gamma (GMG) distribution has been defined. Some important properties of this newly defined distribution are given in Section 4. In Section 5, using the conditioning approach for the matrix variate normal distribution, a new matrix t type family of distributions is introduced. Some important statistical characteristics of this family are studied in Section 6. A Bayesian application is given in Section 7. The paper is concluded in Section 8.

---

### 3. Generalized matrix gamma distribution

---

Recently, Nagar et al. [30] defined a generalized matrix variate gamma distribution by generalizing the multivariate gamma function. We also refer to

Nagar et al. [29] for further generalizations. In this paper, by incorporating an additional factor in the p.d.f, we give a generalization of the matrix variate gamma distribution (Das and Dey [6], Iranmanesh et al. [17]).

In the following we provide the reader with the definition of the generalized matrix variate gamma distribution.

**Definition 3.1.** A random symmetric matrix  $\mathbf{X}$  of order  $p$  is said to have a generalized matrix gamma (GMG) distribution with parameters  $\alpha$ ,  $\beta$ ,  $k$ ,  $\Sigma$  and  $\mathbf{U}$ , denoted by  $\mathbf{X} \sim GMG_p(\alpha, \beta, k, \Sigma, \mathbf{U})$ , if its p.d.f is given by

$$C(\alpha, \beta, k, \Sigma, \mathbf{U}) \left( -\frac{1}{\beta} \Sigma^{-1} \mathbf{X} \right) \det(\mathbf{X})^{\alpha-(p+1)/2} [\text{tr}(\mathbf{X}\mathbf{U})]^k, \quad \mathbf{X} > 0, \quad (3.1)$$

where  $\alpha > (p-1)/2$ ,  $\beta > 0$ ,  $\Sigma > 0$ ,  $\mathbf{U} > 0$ ,  $k \in \mathbb{N}_0$  and  $C(\alpha, \beta, k, \Sigma, \mathbf{U})$  is the normalizing constant.

By integrating the p.d.f of  $\mathbf{X}$  over its support set, the normalizing constant  $C(\alpha, \beta, k, \Sigma, \mathbf{U})$  can be evaluated as

$$\begin{aligned} [C(\alpha, \beta, k, \Sigma, \mathbf{U})]^{-1} &= \int_{\mathbf{X} > 0} \left( -\frac{1}{\beta} \Sigma^{-1} \mathbf{X} \right) \det(\mathbf{X})^{\alpha-(p+1)/2} [\text{tr}(\mathbf{X}\mathbf{U})]^k d\mathbf{X} \\ &= \beta^{p\alpha+k} \Gamma_p(\alpha) \det(\Sigma)^\alpha \sum_{\kappa} (\alpha)_\kappa C_\kappa(\mathbf{U}\Sigma), \end{aligned} \quad (3.2)$$

where the last line has been obtained by using (2.3).

The distribution given by the p.d.f (3.1) is a generalization of the matrix variate gamma distribution (Das and Dey [6], Iranmanesh et al. [17]). For  $\mathbf{U} = \Sigma^{-1}$ , the p.d.f in (3.1) simplifies to

$$\frac{(-\Sigma^{-1} \mathbf{X} / \beta) \det(\mathbf{X})^{\alpha-(p+1)/2} [\text{tr}(\Sigma^{-1} \mathbf{X})]^k}{\beta^{\alpha p+k} (\alpha p)_k \Gamma_p(\alpha) \det(\Sigma)^\alpha}, \quad \mathbf{X} > 0. \quad (3.3)$$

Further, for  $\mathbf{U} = 0$  or  $k = 0$  the p.d.f (3.1) reduces to the matrix variate gamma p.d.f given by

$$\frac{(-\Sigma^{-1} \mathbf{X} / \beta) \det(\mathbf{X})^{\alpha-(p+1)/2}}{\beta^{\alpha p} \Gamma_p(\alpha) \det(\Sigma)^\alpha}, \quad \mathbf{X} > 0. \quad (3.4)$$

By suitably choosing  $\beta$  we can derive a number of special cases of (3.3). If we choose  $\alpha = n/2$  and  $\beta = 2$ , then  $\mathbf{X}$  has a generalized Wishart distribution with p.d.f

$$\frac{(-\Sigma^{-1} \mathbf{X} / 2) \det(\mathbf{X})^{n/2-(p+1)/2} [\text{tr}(\Sigma^{-1} \mathbf{X})]^k}{2^{np/2+k} \Gamma_p(n/2) (np/2)_k \det(\Sigma)^{n/2}}, \quad \mathbf{X} > 0. \quad (3.5)$$

Note that  $n$  is a positive integer, generally considered as the sample size. If we choose  $\Sigma = \mathbf{I}_p$ ,  $\beta = 2$  and  $p = 1$  in (3.3), then the scalar variable  $X$  follows

a chi-square distribution with  $n + 2k$  degrees of freedom. Further, if we take  $p = 1$  and  $\beta = 1$  in (3.3), then the scalar variable  $X$  follows a univariate gamma distribution with shape parameter  $\alpha + k$  and scale parameter  $\sigma$ . Finally, for  $\Sigma = \mathbf{I}_p$  and  $p = 1$ , the scalar variable  $X$  follows a univariate gamma distribution with shape parameter  $\alpha + k$  and scale parameter  $\beta$ .

**Definition 3.2.** red If  $\mathbf{X} \sim GMG_p(\alpha, \beta, k, \Sigma, \mathbf{U})$  then  $\mathbf{X}^{-1}$  is said to have an inverted generalized matrix gamma (IGMG) distribution with parameters  $\alpha, \beta, k, \Sigma^{-1}$  and  $\mathbf{U}$ , denoted by  $\mathbf{X}^{-1} \sim IGMG_p(\alpha, \beta, k, \Sigma^{-1}, \mathbf{U})$ .

red In the following theorem, the p.d.f. of the IGMG distribution is derived.

**Proposition 3.1.** red Let  $\mathbf{X} \sim GMG_p(\alpha, \beta, k, \Sigma, \mathbf{U})$ . Then,  $\mathbf{Y} = \mathbf{X}^{-1} \sim IGMG_p(\alpha, \beta, k, \Sigma^{-1}, \mathbf{U})$  has the p.d.f given by

$$(\mathfrak{C}(\alpha, \beta, k, \Sigma^{-1}, \mathbf{U}) \left( -\frac{1}{\beta} \Sigma \mathbf{Y}^{-1} \right) \det(\mathbf{Y})^{-\alpha-(p+1)/2} [\text{tr}(\mathbf{Y}^{-1} \mathbf{U})]^k, \quad \mathbf{Y} > 0,$$

where  $\alpha > (p-1)/2$ ,  $\beta > 0$ ,  $\Sigma > 0$ ,  $\mathbf{U} > 0$ ,  $k \in \mathbb{N}_0$  and  $C(\alpha, \beta, k, \Sigma^{-1}, \mathbf{U})$  is the normalizing constant.

**Proof:** The proof follows from the fact that the Jacobian of the transformation  $\mathbf{Y} = \mathbf{X}^{-1}$  is given by  $J(\mathbf{X} \rightarrow \mathbf{Y}) = \det(\mathbf{Y})^{-(p+1)}$ .  $\square$

red By taking  $\mathbf{U} = \Sigma$ ,  $\alpha = n/2$  and  $\beta = 2$  in (3.6), the inverted generalized Wishart p.d.f can be obtained as

$$(3.7) \quad \frac{(-\Sigma \mathbf{Y}^{-1}/2) \det(\mathbf{Y})^{-n/2-(p+1)/2} [\text{tr}(\Sigma \mathbf{Y}^{-1})]^k}{2^{np/2+k} \Gamma_p(n/2) (np/2)_k \det(\Sigma)^{-n/2}}, \quad \mathbf{Y} > 0.$$

---

#### 4. Properties of GMG and IGMG distributions

---

In this section, various properties of the GMG and IGMG distributions are derived.

**Proposition 4.1.** Let  $\mathbf{X} \sim GMG_p(\alpha, \beta, k, \Sigma, \mathbf{U})$ . Then, the Laplace transform of  $\mathbf{X}$  is

$$\varphi_{\mathbf{X}}(\mathbf{T}) = \det(\mathbf{I}_p + \beta \Sigma \mathbf{T})^{-\alpha} \frac{\sum_{\kappa} (\alpha)_{\kappa} C_{\kappa} \left( \mathbf{U} (\beta \mathbf{T} + \Sigma^{-1})^{-1} \right)}{\sum_{\kappa} (\alpha)_{\kappa} C_{\kappa} (\mathbf{U} \Sigma)}, \quad (4.1)$$

where  $\mathbf{T}$  is a complex symmetric matrix of order  $p$  with  $\text{Re}(\mathbf{T}) > 0$ .



**Proof:** By definition, we have

$$\begin{aligned}\varphi_{\mathbf{X}}(\mathbf{T}) &= E[\exp(-\text{tr}(\mathbf{T}\mathbf{X}))] \\ &= C(\alpha, \beta, k, \mathbf{\Sigma}, \mathbf{U}) \int_{\mathbf{X} > 0} \left[ -\mathbf{X}(\mathbf{T} + \frac{1}{\beta}\mathbf{\Sigma}^{-1}) \right] \det(\mathbf{X})^{\alpha-(p+1)/2} \\ &\quad \times [\text{tr}(\mathbf{X}\mathbf{U})]^k d\mathbf{X}.\end{aligned}$$

Now, evaluating the above integral by using (3.2) and simplifying the resulting expression, we get the desired result.  $\square$

**Corollary 4.0.1.** Let  $\mathbf{X} \sim GMG_p(\alpha, \beta, k, \mathbf{\Sigma}, \mathbf{U})$ . Then the characteristic function of  $\mathbf{X}$  is

$$\psi_{\mathbf{X}}(\mathbf{T}) = \det(\mathbf{I}_p - \iota\beta\mathbf{\Sigma}\mathbf{T})^{-\alpha} \frac{\sum_{\kappa}(\alpha)_{\kappa} C_{\kappa}(\mathbf{U}\mathbf{\Sigma}(\mathbf{I}_p - \iota\beta\mathbf{T}\mathbf{\Sigma})^{-1})}{\sum_{\kappa}(\alpha)_{\kappa} C_{\kappa}(\mathbf{U}\mathbf{\Sigma})}, \quad (4.2)$$

where  $\iota = \sqrt{-1}$ ,  $\mathbf{T}$  is a symmetric positive definite matrix of order  $p$  with  $\mathbf{T} = ((1 + \delta_{ij})t_{ij}/2)$  and  $\delta_{ij}$  is the Kronecker delta.

**Proposition 4.2.** If  $\mathbf{X} \sim GMG_p(\alpha, \beta, k, \mathbf{\Sigma}, \mathbf{U})$ , then for a  $p \times p$  non-singular constant matrix  $\mathbf{A}$ , we have

$$\mathbf{A}\mathbf{X}\mathbf{A}' \sim GMG_p(\alpha, \beta, k, \mathbf{A}\mathbf{\Sigma}\mathbf{A}', (\mathbf{A}^{-1})'\mathbf{U}\mathbf{A}^{-1}).$$

**Proposition 4.3.** Let  $\mathbf{X} \sim GMG_p(\alpha, \beta, k, \mathbf{\Sigma}, \mathbf{U})$ . Then

$$E[\det(\mathbf{X})^h] = \det(\beta\mathbf{\Sigma})^h \frac{\Gamma_p(\alpha + h)}{\Gamma_p(\alpha)} \frac{\sum_{\kappa}(\alpha + h)_{\kappa} C_{\kappa}(\mathbf{U}\mathbf{\Sigma})}{\sum_{\kappa}(\alpha)_{\kappa} C_{\kappa}(\mathbf{U}\mathbf{\Sigma})}.$$

**Proof:** By definition

$$\begin{aligned}E[\det(\mathbf{X})^h] &= C(\alpha, \beta, k, \mathbf{\Sigma}, \mathbf{U}) \int_{\mathbf{X} > 0} (-\frac{1}{\beta}\mathbf{\Sigma}^{-1}\mathbf{X}) \det(\mathbf{X})^{\alpha+h-(p+1)/2} \\ &\quad \times [\text{tr}(\mathbf{X}\mathbf{U})]^k d\mathbf{X} \\ &= \frac{C(\alpha, \beta, k, \mathbf{\Sigma}, \mathbf{U})}{C(\alpha + h, \beta, k, \mathbf{\Sigma}, \mathbf{U})}.\end{aligned}$$

Now, simplification of the above expression yields the desired result.  $\square$

**Proposition 4.4.** If  $\mathbf{X} \sim GMG_p(\alpha, \beta, k, \mathbf{\Sigma}, \mathbf{\Sigma}^{-1})$ . Then

$$E[\det(\mathbf{X})^h] = \det(\beta\mathbf{\Sigma})^h \frac{\Gamma_p(\alpha + h)}{\Gamma_p(\alpha)} \frac{(\alpha p + hp)_k}{(\alpha p)_k}.$$

In order to find the expectation of the trace of a GMG random matrix, it is useful to find the expectation of zonal polynomials, which is derived below.

**Theorem 4.1.** *Let  $\mathbf{X} \sim GMG_p(\alpha, \beta, k, \mathbf{\Sigma}, \mathbf{U})$  and  $\mathbf{B}$  be a constant symmetric matrix of order  $p$ . Then*

$$E[C_\tau(\mathbf{XB})] = C(\alpha, \beta, k, \mathbf{\Sigma}, \mathbf{U}) \beta^{p\alpha+t+k} \det(\mathbf{\Sigma})^\alpha \\ \times \sum_{\kappa} \sum_{\phi \in \kappa \cdot \tau} \theta_\phi^{\kappa, \tau} \Gamma_p(\alpha, \phi) C_\phi^{\kappa, \tau}(\mathbf{U}\mathbf{\Sigma}, \mathbf{B}\mathbf{\Sigma}).$$

**Proof:** By definition, we have

$$E[C_\tau(\mathbf{XB})] = C(\alpha, \beta, k, \mathbf{\Sigma}, \mathbf{U}) \\ \times \int_{\mathbf{X} > 0} \left(-\frac{1}{\beta} \mathbf{\Sigma}^{-1} \mathbf{X}\right) \det(\mathbf{X})^{\alpha-(p+1)/2} [\text{tr}(\mathbf{XU})]^k C_\tau(\mathbf{XB}) d\mathbf{X}.$$

Now, writing

$$[\text{tr}(\mathbf{XU})]^k C_\tau(\mathbf{XB}) = \sum_{\kappa} C_\kappa(\mathbf{XU}) C_\tau(\mathbf{XB}) \\ = \sum_{\kappa} \sum_{\phi \in \kappa \cdot \tau} \theta_\phi^{\kappa, \tau} C_\phi^{\kappa, \tau}(\mathbf{XU}, \mathbf{XB}),$$

where we have used (2.7), and integrating  $\mathbf{X}$  by using (2.8), we obtain

$$E[C_\tau(\mathbf{XB})] = C(\alpha, \beta, k, \mathbf{\Sigma}, \mathbf{U}) \sum_{\kappa} \sum_{\phi \in \kappa \cdot \tau} \theta_\phi^{\kappa, \tau} \\ \times \int_{\mathbf{X} > 0} \left(-\frac{1}{\beta} \mathbf{\Sigma}^{-1} \mathbf{X}\right) \det(\mathbf{X})^{\alpha-(p+1)/2} C_\phi^{\kappa, \tau}(\mathbf{XU}, \mathbf{XB}) d\mathbf{X}, \\ = C(\alpha, \beta, k, \mathbf{\Sigma}, \mathbf{U}) \det(\beta\mathbf{\Sigma})^\alpha \sum_{\kappa} \sum_{\phi \in \kappa \cdot \tau} \theta_\phi^{\kappa, \tau} \Gamma_p(\alpha, \phi) C_\phi^{\kappa, \tau}(\beta\mathbf{U}\mathbf{\Sigma}, \beta\mathbf{B}\mathbf{\Sigma}).$$

Now, the result follows from the fact that

$$C_\phi^{\kappa, \tau}(\beta\mathbf{U}\mathbf{\Sigma}, \beta\mathbf{B}\mathbf{\Sigma}) = \beta^{k+t} C_\phi^{\kappa, \tau}(\mathbf{U}\mathbf{\Sigma}, \mathbf{B}\mathbf{\Sigma}).$$

□

**Theorem 4.2.** *Let  $\mathbf{Y} \sim IGMG_p(\alpha, \beta, k, \mathbf{\Psi}, \mathbf{U})$ . Then, the Laplace transform of  $\mathbf{Y}$  is given by*

$$(4.3) \varphi_{\mathbf{Y}}(\mathbf{T}) = C(\alpha, \beta, k, \mathbf{\Psi}^{-1}, \mathbf{U}) \det(\mathbf{T})^\alpha \left[ \frac{d^k}{dz^k} B_\alpha(\mathbf{T}(\beta^{-1}\mathbf{\Psi} - z\mathbf{U})) \right]_{z=0},$$

where  $\mathbf{T}$  is a complex symmetric matrix of order  $p$  with  $\text{Re}(\mathbf{T}) > 0$  and  $B_\delta(\cdot)$  is the Bessel function of matrix argument (Herz [16]) given by

$$B_\delta(\mathbf{WZ}) = \det(\mathbf{W})^{-\delta} \int_{\mathbf{S} > 0} \det(\mathbf{S})^{\delta-(p+1)/2} (-\mathbf{SZ} - \mathbf{S}^{-1}\mathbf{W}) d\mathbf{S}. \quad (4.4)$$

**Proof:** The Laplace transform of  $\mathbf{Y}$ , denoted by  $\varphi_{\mathbf{Y}}(\mathbf{T})$  can be derived as

$$\begin{aligned} \varphi_{\mathbf{Y}}(\mathbf{T}) &= C(\alpha, \beta, k, \mathbf{\Psi}^{-1}, \mathbf{U}) \\ &\quad \times \int_{\mathbf{Y} > 0} (-\mathbf{T}\mathbf{Y}) \left( -\frac{1}{\beta} \mathbf{\Psi}\mathbf{Y}^{-1} \right) \det(\mathbf{Y})^{-\alpha-(p+1)/2} [\text{tr}(\mathbf{Y}^{-1}\mathbf{U})]^k d\mathbf{Y}. \end{aligned} \quad (4.5)$$

Note that, we can write

$$[\text{tr}(\mathbf{Y}^{-1}\mathbf{U})]^k = \left[ \frac{d^k}{dz^k} \exp[z \text{tr}(\mathbf{Y}^{-1}\mathbf{U})] \right]_{z=0}. \quad (4.6)$$

Now, substituting (4.6) in (4.5), we have

$$\begin{aligned} \varphi_{\mathbf{Y}}(\mathbf{T}) &= C(\alpha, \beta, k, \mathbf{\Psi}^{-1}, \mathbf{U}) \\ &\quad \times \left[ \frac{d^k}{dz^k} \int_{\mathbf{Y} > 0} (-\mathbf{T}\mathbf{Y}) [- (\beta^{-1}\mathbf{\Psi} - z\mathbf{U}) \mathbf{Y}^{-1}] \det(\mathbf{Y})^{-\alpha-(p+1)/2} d\mathbf{Y} \right]_{z=0} \\ &= C(\alpha, \beta, k, \mathbf{\Psi}^{-1}, \mathbf{U}) \\ &\quad \times \left[ \frac{d^k}{dz^k} \int_{\mathbf{Y} > 0} [-\mathbf{T}\mathbf{Y}^{-1} - (\beta^{-1}\mathbf{\Psi} - z\mathbf{U}) \mathbf{Y}] \det(\mathbf{Y})^{\alpha-(p+1)/2} d\mathbf{Y} \right]_{z=0}. \end{aligned} \quad (4.7)$$

Finally, using (4.4) in (4.7), we get the desired result.  $\square$

**Proposition 4.5.** Let  $\mathbf{Y} \sim \text{IGMG}_p(\alpha, \beta, k, \mathbf{\Psi}, \mathbf{U})$ . Then

$$E \left[ \det(\mathbf{Y})^h \right] = \frac{\det(\mathbf{\Psi})^h \Gamma_p(\alpha - h) \sum_{\kappa} (\alpha - h)_{\kappa} C_{\kappa}(\mathbf{U}\mathbf{\Psi}^{-1})}{\beta^{ph} \Gamma_p(\alpha) \sum_{\kappa} (\alpha)_{\kappa} C_{\kappa}(\mathbf{U}\mathbf{\Psi}^{-1})}, \quad \text{Re}(\alpha - h) > \frac{p-1}{2}.$$

**Proof:** By definition,

$$\begin{aligned} E \left[ \det(\mathbf{Y})^h \right] &= C(\alpha, \beta, k, \mathbf{\Psi}^{-1}, \mathbf{U}) \\ &\quad \times \int_{\mathbf{Y} > 0} (-\beta^{-1}\mathbf{\Psi}\mathbf{Y}^{-1}) \det(\mathbf{Y})^{-(\alpha-h)-(p+1)/2} [\text{tr}(\mathbf{Y}^{-1}\mathbf{U})]^k d\mathbf{Y} \\ &= \frac{C(\alpha, \beta, k, \mathbf{\Psi}^{-1}, \mathbf{U})}{C(\alpha - h, \beta, k, \mathbf{\Psi}^{-1}, \mathbf{U})}, \quad \text{Re}(\alpha - h) > \frac{p-1}{2}. \end{aligned}$$

Now, the desired result is obtained by simplifying the above expression.  $\square$

**Proposition 4.6.** Let  $\mathbf{Y} \sim \text{IGMG}_p(\alpha, \beta, k, \mathbf{\Psi}, \mathbf{U})$  and  $\mathbf{A}$  be a constant symmetric matrix of order  $p$ . Then  $\mathbf{A}\mathbf{Y}\mathbf{A}' \sim \text{IGMG}_p(\alpha, \beta, k, \mathbf{A}\mathbf{\Psi}\mathbf{A}', \mathbf{A}\mathbf{U}\mathbf{A}')$ .

**Proof:** The Jacobian of the transformation  $\mathbf{Z} = \mathbf{A}\mathbf{Y}\mathbf{A}'$  is  $J(\mathbf{Y} \rightarrow \mathbf{Z}) = \det(\mathbf{A})^{-(p+1)}$ . Substituting appropriately in the p.d.f of  $\mathbf{Y}$ , we get the desired result.  $\square$

**Theorem 4.3.** Let the  $p \times p$  random symmetric matrices  $\mathbf{X}_1$  and  $\mathbf{X}_2$  be independent,  $\mathbf{X}_1 \sim \text{GMG}_p(\alpha_1, \beta, k, \boldsymbol{\Sigma}, \mathbf{U})$  and  $\mathbf{X}_2 \sim \text{GMG}_p(\alpha_2, \beta, l, \boldsymbol{\Sigma}, \mathbf{U})$ . Define  $\mathbf{R} = (\mathbf{X}_1 + \mathbf{X}_2)^{-1/2} \mathbf{X}_1 (\mathbf{X}_1 + \mathbf{X}_2)^{-1/2}$  and  $\mathbf{S} = \mathbf{X}_1 + \mathbf{X}_2$ . The p.d.f of  $\mathbf{S}$  is given by

$$C(\alpha_1, \beta, k, \boldsymbol{\Sigma}, \mathbf{U}) C(\alpha_2, \beta, l, \boldsymbol{\Sigma}, \mathbf{U}) \left[ -(\beta \boldsymbol{\Sigma})^{-1} \mathbf{S} \right] \det(\mathbf{S})^{\alpha_1 + \alpha_2 - (p+1)/2} \\ \times \sum_{\kappa} \sum_{\lambda} \sum_{\phi \in \kappa \cdot \lambda} \theta_{\phi}^{\kappa, \lambda} \frac{\Gamma_p(\alpha_1, \kappa) \Gamma_p(\alpha_1, \lambda)}{\Gamma_p(\alpha_1 + \alpha_2, \phi)} C_{\phi}^{\kappa, \lambda}(\mathbf{S}\mathbf{U}, \mathbf{S}\mathbf{U}), \quad \mathbf{S} > 0.$$

Further, for  $\mathbf{U} = \mathbf{I}_p$ , the p.d.f of  $\mathbf{R}$  is given by

$$C(\alpha_1, \beta, k, \boldsymbol{\Sigma}, \mathbf{I}_p) C(\alpha_2, \beta, l, \boldsymbol{\Sigma}, \mathbf{I}_p) \det(\beta \boldsymbol{\Sigma})^{\alpha_1 + \alpha_2} \det(\mathbf{R})^{\alpha_1 - (p+1)/2} \det(\mathbf{I}_p - \mathbf{R})^{\alpha_2 - (p+1)/2} \\ \times \sum_{\kappa} \sum_{\lambda} \sum_{\phi \in \kappa \cdot \lambda} \theta_{\phi}^{\kappa, \lambda} \Gamma_p(\alpha_1 + \alpha_2, \phi) C_{\phi}^{\kappa, \lambda}(\beta \boldsymbol{\Sigma} \mathbf{R}, \beta \boldsymbol{\Sigma} (\mathbf{I}_p - \mathbf{R})), \quad 0 < \mathbf{R} < \mathbf{I}_p.$$

**Proof:** The joint p.d.f of  $\mathbf{X}_1$  and  $\mathbf{X}_2$  is given by

$$C(\alpha_1, \beta, k, \boldsymbol{\Sigma}, \mathbf{U}) C(\alpha_2, \beta, l, \boldsymbol{\Sigma}, \mathbf{U}) \left[ -(\beta \boldsymbol{\Sigma})^{-1} (\mathbf{X}_1 + \mathbf{X}_2) \right] \\ \times \det(\mathbf{X}_1)^{\alpha_1 - (p+1)/2} \det(\mathbf{X}_2)^{\alpha_2 - (p+1)/2} [\text{tr}(\mathbf{X}_1 \mathbf{U})]^k [\text{tr}(\mathbf{X}_2 \mathbf{U})]^l, \quad \mathbf{X}_1 > 0, \quad \mathbf{X}_2 > 0.$$

Transforming  $\mathbf{R} = (\mathbf{X}_1 + \mathbf{X}_2)^{-1/2} \mathbf{X}_1 (\mathbf{X}_1 + \mathbf{X}_2)^{-1/2}$  and  $\mathbf{S} = \mathbf{X}_1 + \mathbf{X}_2$  with the Jacobian  $J(\mathbf{X}_1, \mathbf{X}_2 \rightarrow \mathbf{R}, \mathbf{S}) = \det(\mathbf{S})^{(p+1)/2}$  in the joint p.d.f of  $\mathbf{X}_1$  and  $\mathbf{X}_2$ , the joint p.d.f of  $\mathbf{R}$  and  $\mathbf{S}$  can be derived as

$$C(\alpha_1, \beta, k, \boldsymbol{\Sigma}, \mathbf{U}) C(\alpha_2, \beta, l, \boldsymbol{\Sigma}, \mathbf{U}) \left[ -(\beta \boldsymbol{\Sigma})^{-1} \mathbf{S} \right] \det(\mathbf{S})^{\alpha_1 + \alpha_2 - (p+1)/2} \\ \times \det(\mathbf{R})^{\alpha_1 - (p+1)/2} \det(\mathbf{I}_p - \mathbf{R})^{\alpha_2 - (p+1)/2} \left[ \text{tr}(\mathbf{S}^{1/2} \mathbf{R} \mathbf{S}^{1/2} \mathbf{U}) \right]^k \left[ \text{tr}(\mathbf{S}^{1/2} (\mathbf{I}_p - \mathbf{R}) \mathbf{S}^{1/2} \mathbf{U}) \right]^l, \quad (4.8)$$

where  $\mathbf{S} > 0$  and  $0 < \mathbf{R} < \mathbf{I}_p$ . Now, writing

$$\left[ \text{tr}(\mathbf{S}^{1/2} \mathbf{R} \mathbf{S}^{1/2} \mathbf{U}) \right]^k \left[ \text{tr}(\mathbf{S}^{1/2} (\mathbf{I}_p - \mathbf{R}) \mathbf{S}^{1/2} \mathbf{U}) \right]^l \\ = \sum_{\kappa} \sum_{\lambda} \sum_{\phi \in \kappa \cdot \lambda} \theta_{\phi}^{\kappa, \lambda} C_{\phi}^{\kappa, \lambda}(\mathbf{S}^{1/2} \mathbf{U} \mathbf{S}^{1/2} \mathbf{R}, \mathbf{S}^{1/2} \mathbf{U} \mathbf{S}^{1/2} (\mathbf{I}_p - \mathbf{R}))$$

in (4.8), the joint p.d.f of  $\mathbf{R}$  and  $\mathbf{S}$  can be re-written as

$$C(\alpha_1, \beta, k, \boldsymbol{\Sigma}, \mathbf{U}) C(\alpha_2, \beta, l, \boldsymbol{\Sigma}, \mathbf{U}) \left[ -(\beta \boldsymbol{\Sigma})^{-1} \mathbf{S} \right] \det(\mathbf{S})^{\alpha_1 + \alpha_2 - (p+1)/2} \\ \times \det(\mathbf{R})^{\alpha_1 - (p+1)/2} \det(\mathbf{I}_p - \mathbf{R})^{\alpha_2 - (p+1)/2} \\ \times \sum_{\kappa} \sum_{\lambda} \sum_{\phi \in \kappa \cdot \lambda} \theta_{\phi}^{\kappa, \lambda} C_{\phi}^{\kappa, \lambda}(\mathbf{S}^{1/2} \mathbf{U} \mathbf{S}^{1/2} \mathbf{R}, \mathbf{S}^{1/2} \mathbf{U} \mathbf{S}^{1/2} (\mathbf{I}_p - \mathbf{R})), \quad (4.9)$$

where  $\mathbf{S} > 0$  and  $0 < \mathbf{R} < \mathbf{I}_p$ . Finally, integrating the above expression with respect to  $\mathbf{R}$  by using (2.13), we get the p.d.f of  $\mathbf{S}$ . Further, substituting  $\mathbf{U} = \mathbf{I}_p$  in the above expression and integrating  $\mathbf{S}$  by using (2.8), we get the p.d.f of  $\mathbf{R}$ .  $\square$

---

## 5. Family of generalized matrix variate $t$ -distributions

---

In this section, a new family of matrix variate  $t$  distributions is introduced. This distribution will be useful in Bayesian analysis.

**Definition 5.1.** The  $n \times p$  random matrix  $\mathbf{T}$  is said to have a generalized matrix variate  $t$  distribution (GMT) with parameters  $\mathbf{M} \in \mathbb{R}^{n \times p}$ ,  $\Psi(p \times p) > 0$ ,  $\Omega(n \times n) > 0$ ,  $\mathbf{U}(p \times p) > 0$ ,  $\alpha > (p-1)/2$ ,  $\beta > 0$ ,  $\kappa = (k_1, \dots, k_p)$ ,  $k_1 \geq \dots \geq k_p \geq 0$ , if its p.d.f is given by

$$\begin{aligned} & \frac{\det(\Omega)^{-p/2} \det(\Psi)^{-n/2} \Gamma_p(\alpha + n/2)}{\Gamma_p(\alpha) \sum_{\kappa} (\alpha)_{\kappa} C_{\kappa}(\mathbf{U} \Psi^{-1})} \left( \frac{\beta}{2\pi} \right)^{np/2} \\ & \times \det \left( \mathbf{I}_n + \frac{\beta}{2} \Omega^{-1} (\mathbf{T} - \mathbf{M}) \Psi^{-1} (\mathbf{T} - \mathbf{M})' \right)^{-(\alpha + n/2)} \\ & \times \sum_{\kappa} \left( \alpha + \frac{n}{2} \right)_{\kappa} C_{\kappa} \left( \mathbf{U} \left( \Psi + \frac{\beta}{2} (\mathbf{T} - \mathbf{M})' \Omega^{-1} (\mathbf{T} - \mathbf{M}) \right)^{-1} \right), \quad \mathbf{T} \in \mathbb{R}^{n \times p}. \end{aligned} \quad (5.1)$$

We shall use the notation  $\mathbf{T} \sim GMT_{n,p}(\alpha, \beta, k, \mathbf{M}, \Omega, \Psi, \mathbf{U})$ .

For  $\beta = 2$ ,  $\alpha = (m + p - 1)/2$  and  $k = 0$ , the GMT distribution simplifies to the matrix variate  $t$  distribution (Gupta and Nagar [14]). Further, for  $k = 0$ , the GMT simplifies to the generalized matrix variate  $t$  distribution defined by Iranmanesh et al. [19].

**Theorem 5.1.** Let  $\mathbf{X} | \Sigma \sim N_{n,p}(0, \Omega \otimes \Sigma)$  and  $\Sigma \sim IGMG_p(\alpha, \beta, k, \Psi, \mathbf{U})$ . Then,  $\mathbf{X} \sim GMT_{n,p}(\alpha, \beta, k, 0, \Omega, \Psi, \mathbf{U})$ .

**Proof:** Let  $g(\mathbf{X} | \Sigma)$  be the conditional p.d.f of  $\mathbf{X}$  given  $\Sigma$ . Further, let  $h(\Sigma)$  be the marginal p.d.f of  $\Sigma$ . Then, using conditional method, we find the marginal p.d.f of  $\mathbf{X}$  as

$$f(\mathbf{X}) = \int_{\Sigma > 0} g(\mathbf{X} | \Sigma) h(\Sigma) d\Sigma.$$

Now, substituting for  $g(\mathbf{X} | \Sigma)$  and  $h(\Sigma)$  above, we get the marginal p.d.f of  $\mathbf{X}$  as

$$\begin{aligned} f(\mathbf{X}) &= (2\pi)^{-np/2} \det(\Omega)^{-p/2} C(\alpha, \beta, k, \Psi^{-1}, \mathbf{U}) \\ & \times \int_{\Sigma > 0} \left[ -\frac{1}{\beta} \left( \Psi + \frac{\beta}{2} \mathbf{X}' \Omega^{-1} \mathbf{X} \right) \Sigma^{-1} \right] \det(\Sigma)^{-\alpha - (n+p+1)/2} [\text{tr}(\Sigma^{-1} \mathbf{U})]^k d\Sigma. \end{aligned}$$

Further, substituting  $\Sigma^{-1} = \mathbf{Z}$  with the Jacobian  $J(\Sigma \rightarrow \mathbf{Z}) = \det(\mathbf{Z})^{-(p+1)}$  in the above integral and using (3.2), we get

$$f(\mathbf{X}) = (2\pi)^{-np/2} \det(\mathbf{\Omega})^{-p/2} \frac{C(\alpha, \beta, k, \mathbf{\Psi}^{-1}, \mathbf{U})}{C(\alpha + n/2, \beta, k, (\mathbf{\Psi} + \beta \mathbf{X}' \mathbf{\Omega}^{-1} \mathbf{X} / 2)^{-1}, \mathbf{U})}.$$

Finally, simplifying the above expression, we get

$$\begin{aligned} & \frac{\det(\mathbf{\Omega})^{-p/2} \det(\mathbf{\Psi})^{-n/2} \Gamma_p(\alpha + n/2) \left(\frac{\beta}{2\pi}\right)^{np/2}}{\Gamma_p(\alpha) \sum_{\kappa} (\alpha)_{\kappa} C_{\kappa}(\mathbf{U} \mathbf{\Psi}^{-1})} \det\left(\mathbf{I}_n + \frac{\beta}{2} \mathbf{\Omega}^{-1} \mathbf{X} \mathbf{\Psi}^{-1} \mathbf{X}'\right)^{-(\alpha+n/2)} \\ & \times \sum_{\kappa} \left(\alpha + \frac{n}{2}\right)_{\kappa} C_{\kappa}\left(\mathbf{U} \left(\mathbf{\Psi} + \frac{\beta}{2} \mathbf{X}' \mathbf{\Omega}^{-1} \mathbf{X}\right)^{-1}\right), \quad \mathbf{X} \in \mathbb{R}^{n \times p}, \end{aligned}$$

which is the desired result.  $\square$

Next, in Corollary 5.1.1, Corollary 5.1.2 and Theorem 5.2, we give three different variations of the above theorem.

**Corollary 5.1.1.** *Let  $\mathbf{Y}|\Sigma \sim N_{p,n}(0, \Sigma \otimes \mathbf{\Omega})$  and  $\Sigma \sim IGMG_p(\alpha, \beta, k, \mathbf{\Psi}, \mathbf{U})$ . Then, the marginal p.d.f of  $\mathbf{Y}$  is given by*

$$\begin{aligned} & \frac{\det(\mathbf{\Omega})^{-p/2} \det(\mathbf{\Psi})^{-n/2} \Gamma_p(\alpha + n/2) \left(\frac{\beta}{2\pi}\right)^{np/2}}{\Gamma_p(\alpha) \sum_{\kappa} (\alpha)_{\kappa} C_{\kappa}(\mathbf{U} \mathbf{\Psi}^{-1})} \det\left(\mathbf{I}_p + \frac{\beta}{2} \mathbf{\Psi}^{-1} \mathbf{Y} \mathbf{\Omega}^{-1} \mathbf{Y}'\right)^{-(\alpha+n/2)} \\ & \times \sum_{\kappa} \left(\alpha + \frac{n}{2}\right)_{\kappa} C_{\kappa}\left(\mathbf{U} \left(\mathbf{\Psi} + \frac{\beta}{2} \mathbf{Y} \mathbf{\Omega}^{-1} \mathbf{Y}'\right)^{-1}\right), \quad \mathbf{Y} \in \mathbb{R}^{p \times n}. \end{aligned}$$

**Proof:** Take  $\mathbf{Y} = \mathbf{X}'$  in Theorem 5.1.  $\square$

**Corollary 5.1.2.** *Let  $\mathbf{X}|\mathbf{\Omega} \sim N_{n,p}(0, \mathbf{\Omega} \otimes \Sigma)$  and  $\mathbf{\Omega} \sim IGMG_n(\alpha, \beta, k, \mathbf{\Psi}, \mathbf{U})$ . Then, the marginal p.d.f of  $\mathbf{X}$  is*

$$\begin{aligned} & \frac{\det(\Sigma)^{-n/2} \det(\mathbf{\Psi})^{-p/2} \Gamma_n(\alpha + p/2) \left(\frac{\beta}{2\pi}\right)^{np/2}}{\Gamma_n(\alpha) \sum_{\kappa} (\alpha)_{\kappa} C_{\kappa}(\mathbf{U} \mathbf{\Psi}^{-1})} \det\left(\mathbf{I}_n + \frac{\beta}{2} \mathbf{\Psi}^{-1} \mathbf{X} \Sigma^{-1} \mathbf{X}'\right)^{-(\alpha+p/2)} \\ & \times \sum_{\kappa} \left(\alpha + \frac{p}{2}\right)_{\kappa} C_{\kappa}\left(\mathbf{U} \left(\mathbf{\Psi} + \frac{\beta}{2} \mathbf{X} \Sigma^{-1} \mathbf{X}'\right)^{-1}\right), \quad \mathbf{X} \in \mathbb{R}^{n \times p}. \end{aligned}$$

**Proof:** This result can be obtained from Corollary 5.1.1.  $\square$

**Theorem 5.2.** *Let  $\mathbf{Y}|\mathbf{\Omega} \sim N_{p,n}(0, \Sigma \otimes \mathbf{\Omega})$  and  $\mathbf{\Omega} \sim IGMG_n(\alpha, \beta, k, \mathbf{\Psi}, \mathbf{U})$ . Then,  $\mathbf{Y} \sim GMT_{p,n}(\alpha, \beta, k, 0, \Sigma, \mathbf{\Psi}, \mathbf{U})$ .*

**Proof:** Let  $g(\mathbf{Y}|\mathbf{\Omega})$  be the conditional p.d.f of  $\mathbf{Y}$  given  $\mathbf{\Omega}$ . Further, let  $h(\mathbf{\Omega})$  be the marginal p.d.f of  $\mathbf{\Omega}$ . Then, using conditional method, we find the marginal p.d.f of  $\mathbf{Y}$  as

$$f_{\mathbf{Y}}(\mathbf{Y}) = \int_{\mathbf{\Omega} > 0} g(\mathbf{Y}|\mathbf{\Omega}) h(\mathbf{\Omega}) d\mathbf{\Omega}.$$

Now, substituting for  $g(\mathbf{Y}|\mathbf{\Omega})$  and  $h(\mathbf{\Omega})$  above, we get the marginal p.d.f of  $\mathbf{Y}$  as

$$f_{\mathbf{Y}}(\mathbf{Y}) = (2\pi)^{-np/2} \det(\mathbf{\Sigma})^{-n/2} C(\alpha, \beta, k, \mathbf{\Psi}^{-1}, \mathbf{U}) \\ \times \int_{\mathbf{\Omega} > 0} \left[ -\frac{1}{\beta} \left( \mathbf{\Psi} + \frac{\beta}{2} \mathbf{Y}' \mathbf{\Sigma}^{-1} \mathbf{Y} \right) \mathbf{\Omega}^{-1} \right] \det(\mathbf{\Omega})^{-\alpha - (n+p+1)/2} [\text{tr}(\mathbf{\Omega}^{-1} \mathbf{U})]^k d\mathbf{\Omega}.$$

Further, substituting  $\mathbf{\Omega}^{-1} = \mathbf{Z}$  with the Jacobian  $J(\mathbf{\Omega} \rightarrow \mathbf{Z}) = \det(\mathbf{Z})^{-(p+1)}$  in the above integral and using (3.2), we get

$$f_{\mathbf{Y}}(\mathbf{Y}) = (2\pi)^{-np/2} \det(\mathbf{\Sigma})^{-n/2} \frac{C(\alpha, \beta, k, \mathbf{\Psi}^{-1}, \mathbf{U})}{C(\alpha + p/2, \beta, k, (\mathbf{\Psi} + \beta \mathbf{Y}' \mathbf{\Sigma}^{-1} \mathbf{Y} / 2)^{-1}, \mathbf{U})}.$$

Finally, simplifying the above expression, we get the desired result.  $\square$

---

## 6. Some properties of the GMT family of distributions

---

In this section, various properties of the GMT distribution are derived.

**Proposition 6.1.** Let  $\mathbf{T} \sim \text{GMT}_{n,p}(\alpha, \beta, k, \mathbf{M}, \mathbf{\Omega}, \mathbf{\Psi}, \mathbf{U})$ . Let  $\mathbf{A}(n \times n)$  and  $\mathbf{B}(p \times p)$  be constant nonsingular matrices. Then,  $\mathbf{ATB} \sim \text{GMT}_{n,p}(\alpha, \beta, k, \mathbf{AMB}, \mathbf{A\Omega A'}, \mathbf{B'\Psi B}, \mathbf{B'UB})$ .

**Proof:** Transforming  $\mathbf{W} = \mathbf{ATB}$ , with the Jacobian  $J(\mathbf{T} \rightarrow \mathbf{W}) = \det(\mathbf{A})^{-p} \det(\mathbf{B})^{-n}$ , in the p.d.f (5.1) of  $\mathbf{T}$ , and simplifying the resulting expression, we get the result.  $\square$

**Corollary 6.0.1.** If  $\mathbf{T} \sim \text{GMT}_{n,p}(\alpha, \beta, k, \mathbf{M}, \mathbf{\Omega}, \mathbf{\Psi}, \mathbf{U})$ , then

$$\mathbf{\Omega}^{-1/2} \mathbf{T} \mathbf{B} \sim \text{GMT}_{n,p}(\alpha, \beta, k, \mathbf{\Omega}^{-1/2} \mathbf{M} \mathbf{B}, \mathbf{I}_n, \mathbf{B' \Psi B}, \mathbf{B' U B}),$$

$$\mathbf{AT} \mathbf{\Psi}^{-1/2} \sim \text{GMT}_{n,p}(\alpha, \beta, k, \mathbf{AM} \mathbf{\Psi}^{-1/2}, \mathbf{A\Omega A'}, \mathbf{I}_p, \mathbf{\Psi}^{-1/2} \mathbf{U} \mathbf{\Psi}^{-1/2})$$

and

$$\mathbf{\Omega}^{-1/2} \mathbf{T} \mathbf{\Psi}^{-1/2} \sim \text{GMT}_{n,p}(\alpha, \beta, k, \mathbf{\Omega}^{-1/2} \mathbf{M} \mathbf{\Psi}^{-1/2}, \mathbf{I}_n, \mathbf{I}_p, \mathbf{\Psi}^{-1/2} \mathbf{U} \mathbf{\Psi}^{-1/2}).$$

**Proposition 6.2.** If  $\mathbf{T} \sim GMT_{n,p}(\alpha, \beta, k, \mathbf{M}, \mathbf{\Omega}, \mathbf{\Psi}, \mathbf{U})$ , then for  $n \geq p$ , the p.d.f of  $\mathbf{Z} = (\mathbf{T} - \mathbf{M})' \mathbf{\Omega}^{-1} (\mathbf{T} - \mathbf{M})$  is given by

$$\begin{aligned} & \frac{\det(\mathbf{\Omega})^{-p/2} \det(\mathbf{\Psi})^{-n/2} \Gamma_p(\alpha + n/2)}{\Gamma_p(n/2) \Gamma_p(\alpha) \sum_{\kappa} (\alpha)_{\kappa} C_{\kappa}(\mathbf{U} \mathbf{\Psi}^{-1})} \left( \frac{\beta}{2} \right)^{np/2} \det(\mathbf{Z})^{(n-p-1)/2} \\ & \times \det \left( \mathbf{I}_p + \frac{\beta}{2} \mathbf{\Psi}^{-1} \mathbf{Z} \right)^{-(\alpha+n/2)} \sum_{\kappa} \left( \alpha + \frac{n}{2} \right)_{\kappa} C_{\kappa} \left( \mathbf{U} \left( \mathbf{\Psi} + \frac{\beta}{2} \mathbf{Z} \right)^{-1} \right), \quad \mathbf{Z} > 0. \end{aligned} \quad (6.1)$$

**Proof:** The p.d.f of  $\mathbf{Z}$  is given by

$$\begin{aligned} & \frac{\det(\mathbf{\Omega})^{-p/2} \det(\mathbf{\Psi})^{-n/2} \Gamma_p(\alpha + n/2)}{\Gamma_p(\alpha) \sum_{\kappa} (\alpha)_{\kappa} C_{\kappa}(\mathbf{U} \mathbf{\Psi}^{-1})} \left( \frac{\beta}{2\pi} \right)^{np/2} \sum_{\kappa} \left( \alpha + \frac{n}{2} \right)_{\kappa} \\ & \times \int_{(\mathbf{T}-\mathbf{M})' \mathbf{\Omega}^{-1} (\mathbf{T}-\mathbf{M}) = \mathbf{Z}} \det \left( \mathbf{I}_p + \frac{\beta}{2} \mathbf{\Psi}^{-1} (\mathbf{T} - \mathbf{M})' \mathbf{\Omega}^{-1} (\mathbf{T} - \mathbf{M}) \right)^{-(\alpha+n/2)} \\ & \times C_{\kappa} \left( \mathbf{U} \left( \mathbf{\Psi} + \frac{\beta}{2} (\mathbf{T} - \mathbf{M})' \mathbf{\Omega}^{-1} (\mathbf{T} - \mathbf{M}) \right)^{-1} \right) d\mathbf{Z}, \quad \mathbf{Z} > 0. \end{aligned}$$

Now, evaluating the above integral by using Theorem 1.4.10 of Gupta and Nagar [14], we get the result.  $\square$

The following result is a generalization of the work of Dickey [12].

**Theorem 6.1.** Let  $\mathbf{X} \sim N_{n,p}(0, \mathbf{\Omega} \otimes \mathbf{I}_p)$ , independent of  $\mathbf{S} \sim GMG_p(\alpha, \beta, k, \mathbf{\Lambda}^{-1}, \mathbf{U})$ . Define  $\mathbf{T} = \mathbf{X} \mathbf{S}^{-1/2} + \mathbf{M}$ , where  $\mathbf{M}$  is an  $n \times p$  constant matrix and  $\mathbf{S}^{1/2} (\mathbf{S}^{1/2})' = \mathbf{S}$ . Then, the p.d.f of  $\mathbf{T}$  is given by

$$\begin{aligned} & \frac{\det(\mathbf{\Omega})^{-p/2} \det(\mathbf{\Lambda})^{-n/2} \Gamma_p(\alpha + n/2)}{\Gamma_p(\alpha) \sum_{\kappa} (\alpha)_{\kappa} C_{\kappa}(\mathbf{U} \mathbf{\Lambda}^{-1})} \left( \frac{\beta}{2\pi} \right)^{np/2} \\ & \times \det \left( \mathbf{I}_p + \frac{\beta}{2} \mathbf{\Lambda}^{-1} (\mathbf{T} - \mathbf{M})' \mathbf{\Omega}^{-1} (\mathbf{T} - \mathbf{M}) \right)^{-(\alpha+n/2)} \\ & \times \sum_{\kappa} \left( \alpha + \frac{n}{2} \right)_{\kappa} C_{\kappa} \left( \mathbf{U} \left( \mathbf{\Lambda} + \frac{\beta}{2} (\mathbf{T} - \mathbf{M})' \mathbf{\Omega}^{-1} (\mathbf{T} - \mathbf{M}) \right)^{-1} \right), \quad \mathbf{T} \in \mathbb{R}^{n \times p}. \end{aligned}$$

**Proof:** The joint p.d.f of  $\mathbf{X}$  and  $\mathbf{S}$  is given by

$$\begin{aligned} & \frac{(2\pi)^{-np/2} \det(\mathbf{\Omega})^{-p/2} \det(\mathbf{\Lambda})^{\alpha}}{\beta^{p\alpha+k} \Gamma_p(\alpha) \sum_{\kappa} (\alpha)_{\kappa} C_{\kappa}(\mathbf{U} \mathbf{\Lambda}^{-1})} \det(\mathbf{S})^{\alpha-(p+1)/2} [\text{tr}(\mathbf{S} \mathbf{U})]^k \\ & \times \exp \left[ -\text{tr} \left( \frac{1}{\beta} \mathbf{\Lambda} \mathbf{S} + \frac{1}{2} \mathbf{X}' \mathbf{\Omega}^{-1} \mathbf{X} \right) \right], \quad \mathbf{S} > 0, \quad \mathbf{X} \in \mathbb{R}^{n \times p}. \end{aligned}$$

Now, let  $\mathbf{T} = \mathbf{X} \mathbf{S}^{-1/2} + \mathbf{M}$ . The Jacobian of this transformation is  $J(\mathbf{X} \rightarrow \mathbf{T}) = \det(\mathbf{S})^{n/2}$ . Substituting for  $\mathbf{X}$  in terms of  $\mathbf{T}$  in the joint p.d.f of  $\mathbf{X}$  and  $\mathbf{S}$ ,



and multiplying the resulting expression by  $J(\mathbf{X} \rightarrow \mathbf{T})$ , we get the joint p.d.f of  $\mathbf{T}$  and  $\mathbf{S}$  as

$$\frac{(2\pi)^{-np/2} \det(\mathbf{\Omega})^{-p/2} \det(\mathbf{\Lambda})^\alpha}{\beta^{p\alpha+k} \Gamma_p(\alpha) \sum_{\kappa} (\alpha)_{\kappa} C_{\kappa}(\mathbf{U} \mathbf{\Lambda}^{-1})} \det(\mathbf{S})^{\alpha+n/2-(p+1)/2} [\text{tr}(\mathbf{S} \mathbf{U})]^k \\ \times \left[ -\frac{1}{\beta} \left( \mathbf{\Lambda} + \frac{\beta}{2} (\mathbf{T} - \mathbf{M})' \mathbf{\Omega}^{-1} (\mathbf{T} - \mathbf{M}) \right) \mathbf{S} \right], \quad \mathbf{S} > 0, \quad \mathbf{T} \in \mathbb{R}^{n \times p}.$$

Now, integrating out  $\mathbf{S}$  by using (3.2) and simplifying the resulting expression the p.d.f of  $\mathbf{T}$  is obtained.  $\square$

**Theorem 6.2.** Let  $\mathbf{X} \sim N_{n,p}(0, \mathbf{I}_n \otimes \mathbf{\Sigma})$ , independent of  $\mathbf{S} \sim GMG_n(\alpha, \beta, k, \mathbf{\Lambda}^{-1}, \mathbf{U})$ . Define  $\mathbf{T} = (\mathbf{S}^{-1/2})' \mathbf{X} + \mathbf{M}$ , where  $\mathbf{M}$  is an  $n \times p$  constant matrix and  $\mathbf{S}^{1/2} (\mathbf{S}^{1/2})' = \mathbf{S}$ . Then, the p.d.f of  $\mathbf{T}$  is

$$\frac{\det(\mathbf{\Sigma})^{-n/2} \det(\mathbf{\Lambda})^{-p/2} \Gamma_n(\alpha + p/2)}{\Gamma_n(\alpha) \sum_{\kappa} (\alpha)_{\kappa} C_{\kappa}(\mathbf{U} \mathbf{\Lambda}^{-1})} \left( \frac{\beta}{2\pi} \right)^{np/2} \\ \times \det \left( \mathbf{I}_n + \frac{\beta}{2} \mathbf{\Lambda}^{-1} (\mathbf{T} - \mathbf{M}) \mathbf{\Sigma}^{-1} (\mathbf{T} - \mathbf{M})' \right)^{-(\alpha+p/2)} \\ \times \sum_{\kappa} \left( \alpha + \frac{p}{2} \right)_{\kappa} C_{\kappa} \left( \mathbf{U} \left( \mathbf{\Lambda} + \frac{\beta}{2} (\mathbf{T} - \mathbf{M}) \mathbf{\Sigma}^{-1} (\mathbf{T} - \mathbf{M})' \right)^{-1} \right), \quad \mathbf{T} \in \mathbb{R}^{n \times p}.$$

**Proof:** The joint p.d.f of  $\mathbf{X}$  and  $\mathbf{S}$  is given by

$$\frac{(2\pi)^{-np/2} \det(\mathbf{\Sigma})^{-n/2} \det(\mathbf{\Lambda})^\alpha}{\beta^{n\alpha+k} \Gamma_n(\alpha) \sum_{\kappa} (\alpha)_{\kappa} C_{\kappa}(\mathbf{U} \mathbf{\Lambda}^{-1})} \det(\mathbf{S})^{\alpha-(n+1)/2} [\text{tr}(\mathbf{S} \mathbf{U})]^k \\ \times \exp \left[ -\text{tr} \left( \frac{1}{\beta} \mathbf{\Lambda} \mathbf{S} + \frac{1}{2} \mathbf{X} \mathbf{\Sigma}^{-1} \mathbf{X}' \right) \right], \quad \mathbf{S} > 0, \quad \mathbf{X} \in \mathbb{R}^{n \times p}.$$

Now, let  $\mathbf{T} = (\mathbf{S}^{-1/2})' \mathbf{X} + \mathbf{M}$ . The Jacobian of the transformation is  $J(\mathbf{X} \rightarrow \mathbf{T}) = \det(\mathbf{S})^{p/2}$ . Substituting for  $\mathbf{X}$  in terms of  $\mathbf{T}$  in the joint p.d.f of  $\mathbf{X}$  and  $\mathbf{S}$ , and multiplying the resulting expression by  $J(\mathbf{X} \rightarrow \mathbf{T})$ , we get the joint p.d.f of  $\mathbf{T}$  and  $\mathbf{S}$  as

$$\frac{(2\pi)^{-np/2} \det(\mathbf{\Sigma})^{-n/2} \det(\mathbf{\Lambda})^\alpha}{\beta^{n\alpha+k} \Gamma_n(\alpha) \sum_{\kappa} (\alpha)_{\kappa} C_{\kappa}(\mathbf{U} \mathbf{\Lambda}^{-1})} \det(\mathbf{S})^{\alpha+p/2-(n+1)/2} [\text{tr}(\mathbf{S} \mathbf{U})]^k \\ \times \left[ -\left( \frac{1}{\beta} \mathbf{\Lambda} + \frac{1}{2} (\mathbf{T} - \mathbf{M}) \mathbf{\Sigma}^{-1} (\mathbf{T} - \mathbf{M})' \right) \mathbf{S} \right], \quad \mathbf{S} > 0, \quad \mathbf{T} \in \mathbb{R}^{n \times p}.$$

Now, integrating out  $\mathbf{S}$  by using (3.2) and simplifying the resulting expression, the p.d.f of  $\mathbf{T}$  is obtained.  $\square$

---

## 7. Applications in Bayesian analysis

---

As in Iranmanesh et al. [17], consider the Kullback-Leibler divergence loss (KLDL) function  $\log \left( \frac{\pi(\mathbf{A}|\mathbf{D})}{\pi(\boldsymbol{\Sigma}|\mathbf{D})} \right)$  with the posterior expected loss function

$$\rho(\boldsymbol{\Sigma}, \mathbf{A}) = E \left[ \log \left( \frac{\pi(\mathbf{A}|\mathbf{D})}{\pi(\boldsymbol{\Sigma}|\mathbf{D})} \right) \right].$$

One may use the inverted generalized matrix gamma distribution as a prior distribution in Bayesian context. It is straightforward to prove that posterior distributions are IGMG. They are stated in Propositions 7.1 and 7.2 without proof.

**Proposition 7.1.** *Let  $\mathbf{X}|\boldsymbol{\Sigma} \sim N_{n,p}(0, \boldsymbol{\Omega} \otimes \boldsymbol{\Sigma})$ . Further suppose that the prior distribution of  $\boldsymbol{\Sigma}$  is IGMG with parameters  $(\alpha, \beta, k, \boldsymbol{\Psi}, \mathbf{U})$ . Then, the posterior distribution of  $\boldsymbol{\Sigma}$  is IGMG with parameters  $(\alpha + n/2, \beta, k, (\boldsymbol{\Psi} + \beta \mathbf{X}' \boldsymbol{\Omega}^{-1} \mathbf{X}/2)^{-1}, \mathbf{U})$ . That is, the posterior p.d.f of  $\boldsymbol{\Sigma}$  is*

$$\begin{aligned} \pi(\boldsymbol{\Sigma}|\mathbf{X}) &= C \left( \alpha + \frac{n}{2}, \beta, k, (\boldsymbol{\Psi} + \frac{\beta}{2} \mathbf{X}' \boldsymbol{\Omega}^{-1} \mathbf{X})^{-1}, \mathbf{U} \right) \\ &\times \left[ -\frac{1}{\beta} (\boldsymbol{\Psi} + \frac{\beta}{2} \mathbf{X}' \boldsymbol{\Omega}^{-1} \mathbf{X}) \boldsymbol{\Sigma}^{-1} \right] \det(\boldsymbol{\Sigma})^{-\alpha-(n+p+1)/2} \\ &\times [\text{tr}(\boldsymbol{\Sigma}^{-1} \mathbf{U})]^k, \quad \boldsymbol{\Sigma} > 0. \end{aligned}$$

**Proposition 7.2.** *Let  $\mathbf{X}|\boldsymbol{\Sigma} \sim N_{n,p}(0, \boldsymbol{\Omega} \otimes \boldsymbol{\Sigma})$ . Further suppose that the prior distribution of  $\boldsymbol{\Omega}$  is IGMG with parameters  $(\alpha, \beta, k, \boldsymbol{\Psi}, \mathbf{U})$ . Then, the posterior distribution of  $\boldsymbol{\Omega}$  is IGMG with parameters  $(\alpha + p/2, \beta, k, (\boldsymbol{\Psi} + \beta \mathbf{X} \boldsymbol{\Sigma}^{-1} \mathbf{X}'/2)^{-1}, \mathbf{U})$ . That is, the posterior p.d.f of  $\boldsymbol{\Omega}$  is*

$$\begin{aligned} \pi(\boldsymbol{\Omega}|\mathbf{X}) &= C \left( \alpha + \frac{p}{2}, \beta, k, (\boldsymbol{\Psi} + \frac{\beta}{2} \mathbf{X} \boldsymbol{\Sigma}^{-1} \mathbf{X}')^{-1}, \mathbf{U} \right) \\ &\times \left[ -\frac{1}{\beta} (\boldsymbol{\Psi} + \frac{\beta}{2} \mathbf{X} \boldsymbol{\Sigma}^{-1} \mathbf{X}') \boldsymbol{\Omega}^{-1} \right] \det(\boldsymbol{\Omega})^{-\alpha-(n+p+1)/2} \\ &\times [\text{tr}(\boldsymbol{\Omega}^{-1} \mathbf{U})]^k, \quad \boldsymbol{\Omega} > 0. \end{aligned}$$

By definition, the Bayes estimator of  $\boldsymbol{\Sigma}$ , under the KLDL function, is given by  $\hat{\boldsymbol{\Sigma}} =_{\boldsymbol{\Sigma}} \pi(\boldsymbol{\Sigma}|\mathbf{X})$ . Iranmanesh et al. [19] have shown that

$$\hat{\boldsymbol{\Sigma}} = [\alpha + n/2 + (p+1)/2]^{-1} \left( \frac{1}{2} \mathbf{X}' \boldsymbol{\Omega}^{-1} \mathbf{X} + \frac{1}{\beta} \boldsymbol{\Psi} \right)$$

for the special case  $k = 0$ .

---

## 8. Conclusion

---

In this paper, a generalized matrix variate gamma distribution has been introduced. The corresponding inverted matrix variate gamma distribution has also been derived. By making use of this newly defined matrix variate distribution as the prior for the characteristic matrix of a matrix variate normal distribution, using conditioning approach, a family of generalized matrix variate  $t$  distributions has also been defined.

A future work is to consider estimation of the newly introduced matrix variate distributions. One issue is that the new distributions are over parameterized; that is, there is parameter redundancy. This can be accounted for numerically by constrained maximization of the log likelihood. For example, if the data follow the overparameterized p.d.f.  $ab \exp(-abx)$  then the log likelihood can be maximized using the constraint  $ab = c$ . Usually, partial derivatives of the log likelihood are not required for evaluating maximum likelihood estimates numerically.

---

## ACKNOWLEDGMENTS

---

The authors would like to thank the Editor and the referee for careful reading and comments which greatly improved the paper.

---

## REFERENCES

---

- [1] S. Adhikari, Characterization of uncertainty in damping modeling, In: *Proceedings of the 48th AIAA/ASME/ASCE/AHS/ASC Structures, Structural Dynamics and Materials Conference*, Waikiki, Hawaii (2007).
- [2] redM. Arashi, A. Bekker and J. van Niekerk, (2017) Weighted-type Wishart distributions with application, *REVSTAT-Statistical Journal*, **15** (2017), 205–222.
- [3] A. Bates, A. Daducci and E. Caruyer, Multi-dimensional diffusion MRI sampling scheme: B-tensor design and accurate signal reconstruction, In: *Proceedings of ISMRM 2019 - 27th Annual Meeting Exhibition*, Montreal, Canada (2019), pp. 1–4.
- [4] M. Chamberland, B. Scherrer, S. P. Prabhu, J. Madsen, D. Fortin, K. Whittingstall, M. Descoteaux and S. K. Warfield, Active delineation of Meyer’s loop using oriented priors through MAGNETic tractography (MAGNET), *Human Brain Mapping*, **38** (2017), 509–527.
- [5] Y. Chikuse, Distributions of some matrix variates and latent roots in multivariate Behrens-Fisher discriminant analysis, *Annals of Statistics*, **9** (1981), 401–407.

- [6] S. Das and D. K. Dey, On Bayesian inference for generalized multivariate gamma distribution, *Statistics and Probability Letters*, **80** (2010), 1492–1499.
- [7] S. Das and R. Ghanem, A bounded random matrix approach for stochastic up-scaling, *Multiscale Modeling and Simulation*, **8** (2009), 296–325.
- [8] S. Das and R. Ghanem, Stochastic upscaling for inelastic material behavior from limited experimental data, *Computational Methods for Microstructure-Property Relationships*, editors S. Ghosh and D. Dimiduk, Springer, Boston, MA (2011), pp. 443–468.
- [9] A. W. Davis, Invariant polynomials with two matrix arguments extending the zonal polynomials: Applications to multivariate distribution theory, *Annals of the Institute of Statistical Mathematics*, **31** (1979), no. 4, 465–485.
- [10] A. W. Davis, Invariant polynomials with two matrix arguments extending the zonal polynomials, in: *Multivariate Analysis-V*, editor P. R. Krishnaiah, (1980), pp. 287–299.
- [11] J. A. Díaz-García, Generalisations of some properties of invariant polynomials with matrix arguments, *Applicationes Mathematicae* (Warsaw), **38** (2011), 469–475.
- [12] J. M. Dickey, Matrix variate generalizations of the multivariate  $t$  and the inverted multivariate  $t$  distribution, *Annals of Mathematical Statistics*, **38** (1967), 511–518.
- [13] redJ. T. Ferreira, A. Bekker and M. Arashi, Advances in Wishart type modeling for channel capacity, *REVSTAT-Statistical Journal*, accepted (2018).
- [14] A. K. Gupta and D. K. Nagar, *Matrix Variate Distributions*, Chapman and Hall/CRC, Boca Raton (2000).
- [15] H. J. Haubold and D. Kumar, Fusion yield: Guderley model and Tsallis statistics, *Journal of Plasma Physics*, **77** (2011), 1–14.
- [16] C. S. Herz, Bessel functions of matrix argument, *Annals of Mathematic (2)*, **61** (1955), 474–523.
- [17] A. Iranmanesh, M. Arashi, D. K. Nagar, and S. M. M. Tabatabaey, On inverted matrix variate gamma distribution, *Communications in Statistics - Theory and Methods*, **42** (2013), 28–41.
- [18] A. Iranmanesh, M. Arashi, and S. Shokri, Generalized matrix  $t$  distribution through generalized multivariate gamma distribution, in: *ATINER Conference Paper Series*, No: STA 2014-1114 (2014).
- [19] A. Iranmanesh, M. Arashi and S. M. M. Tabatabaey, On conditional applications of matrix variate normal distribution, *Iranian Journal of Mathematical Sciences and Informatics*, **5** (2010), 33–43.
- [20] A. T. James, Distributions of matrix variate and latent roots derived from normal samples, *Annals of Mathematical Statistics*, **35** (1964), 475–501.
- [21] W. R. Javier and A. K. Gupta, On matrix variate  $t$  distribution, *Communications in Statistics - Theory and Methods*, **14** (1985), 1413–1425.
- [22] B. Jian, B. C. Vemuri, E. Ozarslan, P. Carney and T. Mareci, A continuous mixture of tensors model for diffusion-weighted MR signal reconstruction, In: *Proceedings of the 4th IEEE International Symposium on Biomedical Imaging: From Nano to Macro*, Arlington, VA (2007), pp. 772–775.

- [23] C. G. Khatri, On certain distribution problems based on positive definite quadratic functions in normal vectors, *Annals of Mathematical Statistics*, **37** (1966), 468–479.
- [24] L. Luo, J. Yang, B. Zhang, J. Jiang and H. Huang, Nonparametric Bayesian correlated group regression with applications to image classification, *IEEE Transactions on Neural Networks and Learning Systems*, **29** (2018), 5330–5344.
- [25] D. G. Marx, Quadratic forms of a matrix  $t$  variate, *Annals of the Institute of Statistical Mathematics*, **35** (1983), 347–353.
- [26] A. M. Mathai, *Jacobians of Matrix Transformations and Functions of Matrix Argument*, World Scientific, London (1997).
- [27] R. J. Muirhead, *Aspects of Multivariate Statistical Theory*, John Wiley and Sons, New York (1982).
- [28] D. K. Nagar and A. K. Gupta, Matrix-variate Kummer-beta distribution, *Journal of the Australian Mathematical Society*, **73** (2002), 11–25.
- [29] D. K. Nagar, R. A. Moran-Vasquez, and A. K. Gupta, Extended matrix variate hypergeometric functions and matrix variate distribution, *International Journal of Mathematics and Mathematical Science*, **2015** (2015), doi: 10.1155/2015/190723.
- [30] D. K. Nagar, A. Roldan-Correa, and A. K. Gupta, Extended matrix variate gamma and beta functions, *Journal of Multivariate Analysis*, **122** (2013), 53–69.
- [31] B. Pascual and S. Adhikari, Combined parametricnonparametric uncertainty quantification using random matrix theory and polynomial chaos expansion, *Computers and Structures*, **112** (2012), 364–379.
- [32] B. Scherrer, A. Schwartzman, M. Taquet, S. P. Prabhu, M. Sahin, A. Akhondi-Asl and S. K. Warfield, Characterizing the distribution of anisotropic micro-structural environments with diffusion-weighted imaging (DIAMOND), In: *Medical Image Computing and Computer-Assisted Intervention*, part III, editors K. Mori, I. Sakuma, Y. Sato, C. Barillot and N. Navab, **8151** (2013), pp. 518–526.
- [33] B. Scherrer, M. Taquet, A. Schwartzman, E. St-Onge, G. Rensonnet, S. P. Prabhu and S. K. Warfield, Decoupling axial and radial tissue heterogeneity in diffusion compartment imaging, in: *Information Processing in Medical Imaging*, Lecture Notes in Computer Science, editors M. Niethammer et al., Springer, Cham, **10265** (2017), pp. 440–452.
- [34] A. Takemura, *Zonal Polynomials*, Institute of Mathematical Statistics Lecture Notes Monograph Series, **4**, Institute of Mathematical Statistics, Hayward, CA (1984).